The Evening Sky Map

FREE* EACH MONTH FOR YOU TO EXPLORE, LEARN & ENJOY THE NIGHT SKY

WWW.SKYMAPS.COM

SOUTHERN HEMISPHERE
NOVEMBER 2013

Sky Calendar – November 2013

Get Sky Calendar on Twitter
http://twitter.com/skymaps

1 **Venus at greatest elongation,** 47° east of Sun (evening sky) at 8h UT. Mag. +4.4.

1 **Mercury at inferior conjunction** with the Sun at 20h UT. Mercury passes into the morning sky. Not visible.

3 **New Moon** at 12:48 UT. Start of lunation 1124.

3 **Total Solar Eclipse** from 11:05 to 14:28 UT, greatest eclipse at 12:48 UT. Totality visible along a narrow path between North Atlantic (sunrise) and equatorial Africa (sunset). Partial eclipse in most of Africa, NE South America, southern Europe, and east coast North America.

5 **Moon near Antares** (evening sky) at 12h UT.

6 **Moon at perigee** (closest to Earth) at 9h UT (365,361 km; angular size 32.7').

7 **Moon near Venus** (47° from Sun, evening sky) at 0h UT. Mag. –4.5.

10 **First Quarter Moon** at 5:58 UT.

17 **Full Moon** at 15:15 UT.

17 **Leonid meteor shower peaks** at 10h and 16h UT. Arises from debris ejected by Comet Tempel-Tuttle in 1533. Expect up to 15 meteors per hour under dark skies. Unfavorable this year due to strong moonlight.

18 **Moon near the Pleiades** (midnight sky) at 0h UT.

18 **Mercury at greatest elongation**, 20° west of Sun (morning sky) at 2h UT. Mag. –0.5.

18 **Comet ISON 0.4° N of Spica** (32° from Sun, morning sky) at 2h UT.

18 **Moon near Aldebaran** (morning sky) at 21h UT.

22 **Moon near Jupiter** (morning sky) at 3h UT. Mag. –2.5.

22 **Moon at apogee** (farthest from Earth) at 10h UT (distance 405,443 km; angular size 29.5').

23 **Moon near Beehive Cluster** (morning sky) at 16h UT.

25 **Moon near Regulus** (morning sky) at 13h UT.

25 **Last Quarter Moon** at 19:29 UT.

27 **Moon near Mars** (morning sky) at 13h UT. Mag. +1.3.

28 **Comet ISON at perihelion** at 23h UT.

29 **Moon very near Spica** (morning sky) at 19h UT. Occultation visible in Alaska.

More sky events and links at http://Skymaps.com/skycalendar/

All times in Universal Time (UT). (Australian Eastern Summer Time = UT + 11 hours.)



THIS IS THE BEST TIME OF THE YEAR TO SEE THE GREAT ANDROMEDA GALAXY (M31).

COMPASS DIRECTIONS ARE INDICATED ALONG THE HORIZON CIRCLE (FOR EXAMPLE "NORTH"). TURN THE SKY MAP AROUND ITS CENTER (JUST AS YOU ARE DOING NOW) SO THE COMPASS DIRECTION THAT APPEARS ALONG THE BOTTOM OF THE MAP IS THE SAME AS THE DIRECTION THAT YOU FACE. BEGIN BY USING THE SKY MAP TO FIND A BRIGHT STAR PATTERN IN THE SKY.

THE CELESTIAL OBJECTS ARE LOCATED BETWEEN THE ZENITH AND THE HORIZON.

THE EAST ERN HORIZON CIRCLE IS THE HORIZON.

THE CENTER OF THE MAP IS THE PART OF THE SKY DIRECTLY OVERHEAD (ZENITH) AND THE OUTER EDGE IS THE HORIZON.

SKY MAP SHOWS HOW THE NIGHT SKY LOOKS

EARLY NOV 10 PM
LATE NOV 9 PM
(Add 1 Hour For Daylight Saving)

SKY MAP DRAWN FOR A LATITUDE OF 35° SOUTH AND IS SUITABLE FOR LATITUDES UP TO 15° NORTH OR SOUTH OF THIS

Symbols

- Galaxy
- Double Star
- Variable Star
- Diffuse Nebula
- Planetary Nebula
- Open Star Cluster
- Globular Star Cluster

Star Magnitudes -1 0 1 2 3 4

The constellation Grus represents a water bird, the long-necked Crane.

From southern latitudes stars appear to rotate around the South Celestial Pole (SCP).

INSTRUCTIONS: THE SKY MAP SHOWS THE ENTIRE NIGHT SKY FROM HORIZON-TO-HORIZON AS IT APPEARS ON CERTAIN DATES AND TIMES.

SAVE ON RECOMMENDED PRODUCTS • http://Skymaps.com/store

- Star Atlases & Planispheres
- Star Charts & Astro Posters
- Books for Sky Watchers
- Telescopes & Binoculars

All sales support the production and free distribution of The Evening Sky Map.

Copyright © 2000–2013 Kym Thalassoudis. All Rights Reserved.

* **TERMS OF USE:** FREE FOR NON-COMMERCIAL EDUCATIONAL USE. ASTRONOMY EDUCATION GROUPS MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT http://Skymaps.com/terms.html

About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. **Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

Astronomical Glossary

Conjunction – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

Constellation – A defined area of the sky containing a star pattern.

Diffuse Nebula – A cloud of gas illuminated by nearby stars.

Double Star – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

Elongation – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

Galaxy – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) – The distance a beam of light travels at 300,000 km/sec in one year.

Magnitude – The brightness of a celestial object as it appears in the sky.

Open Star Cluster – A group of tens or hundreds of relatively young stars.

Opposition – When a celestial body is opposite the Sun in the sky.

Planetary Nebula – The remnants of a shell of gas blown off by a star.

Universal Time (UT) – A time system used by astronomers. Also known as Greenwich Mean Time. Australian Eastern Standard Time (Sydney, Australia) is UT plus 10 hours.

Variable Star – A star that changes brightness over a period of time.

SOUTHERN HEMISPHERE
NOVEMBER 2013

CELESTIAL OBJECTS

Skymaps.com

Easily Seen with the Naked Eye

Altair	Aql	•	Brightest star in Aquila. Name means "the flying eagle". Dist=16.7 ly.
Sirius	CMa	•	The brightest star in the sky. Also known as the "Dog Star". Dist=8.6 ly.
Canopus	Car	•	Second brightest star in the sky. 14,000 times more luminous than the Sun. Dist=309 ly.
β Centauri	Cen	•	With Alpha Centauri, forms the so-called "Pointers-to-the-Cross". Dist=525 ly.
α Centauri	Cen	◍	Nearest bright star to Sun at 4.4 ly. Brilliant double star in a telescope. 80 year period.
Achernar	Eri	•	Brightest star in Eridanus, The River. Arabic name meaning "end of river". Dist=140 ly.
Rigel	Ori	•	The brightest star in Orion. Blue supergiant star with mag 7 companion. Dist=770 ly.
Betelgeuse	Ori	•	One of the largest red supergiant stars known. Diameter=300 times that of Sun. Dist=430 ly.
Algol	Per	◉	Famous eclipsing binary star. Magnitude varies between 2.1 & 3.4 over 2.867 days.
Fomalhaut	PsA	•	Brightest star in Piscis Austrinus. In Arabic the "fish's mouth". Dist=25 ly.
Pleiades	Tau	⁘	The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=399 ly.
Hyades	Tau	⁘	Large V-shaped star cluster. Binoculars reveal many more stars. Dist=152 ly.
Aldebaran	Tau	•	Brightest star in Taurus. It is not associated with the Hyades star cluster. Dist=66.7 ly.

Easily Seen with Binoculars

M31	And	⬭	The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.5 million ly.
M2	Aqr	⊕	Resembles a fuzzy star in binoculars.
η Aquilae	Aql	◉	Bright Cepheid variable. Mag varies between 3.6 & 4.5 over 7.166 days. Dist=1,200 ly.
6397	Ara	⊕	Thought to be the nearest globular. Dist=7,000 ly.
M41	CMa	⁘	First recorded observation by Aristotle in 325 BC as "cloudy spot". Dist=2,300 ly.
2516	Car	⁘	Spectacular open star cluster of 100 stars spaning 1/2 deg. Dist=1,300 ly.
2808	Car	⊕	Located 4 deg W of Nu Carinae. Visible to the naked eye on clear nights.
R Carinae	Car	◉	Long period variable. Magnitude varies between 3.9 & 10.5 over 309 days.
Mira	Cet	◉	Famous long period variable star. Mag varies between 3.0 & 10.1 over 332 days.
LMC	Dor	⬭	Large Magellanic Cloud. A neighbouring galaxy of the Milky Way. Dist=180,000 ly.
γ Leporis	Lep	◍	Visible with binoculars. Gold & white stars. Mags 3.6 & 6.2. Dist=30 ly. Sep=96.3".
2232	Mon	⁘	A large scattered star cluster of 20 stars. Dist=1,300 ly.
Cr 69	Ori	⁘	Lambda Orionis Cluster. Dist=1,630 ly.
M42	Ori	□	The Great Orion Nebula. Spectacular bright nebula. Best in telescope. Dist=1,300 light years.
κ Pavonis	Pav	◉	Cepheid-type. Magnitude varies between 3.9 & 4.8 over 9.088 days.
6752	Pav	⊕	One of the better globular star clusters in the sky. Dist=14,000 ly.
M15	Peg	⊕	Only globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly.
ζ Phoenicis	Phe	◍	Eclipsing binary star and double (mag 8). Varies between 3.9 & 4.4 over 1.667 days.
L^2	Pup	◉	Semi-regular variable. Magnitude varies between 2.6 & 6.2 over 140.42 days.
2451	Pup	⁘	30+ stars in binoculars. The brightest star, χ Puppis, is red. Dist=850 ly.
2477	Pup	⁘	Very rich but distant star cluster (4,200 ly). Resembles globular through binoculars.
M25	Sgr	⁘	Bright cluster located about 6 deg N of "teapot's" lid. Dist=1,900 ly.
253	Scl	⬭	Fine, large, cigar-shaped galaxy. Requires dark sky. Member of Sculptor Group.
6025	TrA	⁘	A small open star cluster in Milky Way. Dist=2,700 ly.
47 Tucanae	Tuc	⊕	Spectacular object. Telescope will reveal stars. Near edge of SMC. Dist=15,000 ly.
β Tucanae	Tuc	◍	Complex multiple star. Binoculars show one pair. Telescope required to split primary star.
SMC	Tuc	⬭	Small Magellanic Cloud. Companion galaxy to Milky Way. Requires dark sky. Dist=210,000 ly.
2547	Vel	⁘	Fine open cluster visible through binoculars. Dist=1,300 ly.
IC 2391	Vel	⁘	Omicron Velorum Cluster. Superb object for binoculars. Dist=450 ly.

Telescopic Objects

γ Andromedae	And	◍	Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8".
7009	Aqr	✧	Saturn Nebula. Requires 8-inch telescope to see Saturn-like appendages.
7293	Aqr	✧	Helix Nebula. Spans nearly 1/4 deg. Requires dark sky. Dist=300 ly.
γ Arietis	Ari	◍	Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".
3918	Cen	✧	The Blue Planetary. Visible in a small telescope as a round blue disk.
γ Delphini	Del	◍	Appear yellow & white. Mags 4.3 & 5.2. Dist=100 ly. Struve 2725 double in same field.
2070	Dor	□	Tarantula Nebula. A bright nebula located in LMC. A star-forming region.
θ Eridani	Eri	◍	Striking blue-white double star. Mags 3.2 & 4.3. Visible in a small telescope. Sep=8.2".
β Monocerotis	Mon	◍	Triple star. Mags 4.6, 5.0 & 5.4. Requires telescope to view arc-shape. Sep=7.3".
σ Orionis	Ori	◍	Superb multiple star. 2 mag 7 stars one side, mag 9 star on other. Struve 761 triple in field.
M33	Tri	⬭	Fine face-on spiral galaxy. Requires a large aperture telescope. Dist=2.3 million ly.

The Evening Sky Map (ISSN 1839-7735) Copyright © 2000–2013 Kym Thalassoudis. All Rights Reserved.